arXiv:1401.3119v1 [math.GN] 14 Jan 2014

# ON TOPOLOGICAL STRUCTURES OF FUZZY PARAMETRIZED SOFT SETS

S. ATMACA AND İ. ZORLUTUNA

Abstract. In this paper, we introduce the topological structure of fuzzy parametrized soft sets and fuzzy parametrized soft mappings. We define the notion of quasi-coincidence for fuzzy parametrized soft sets and investigated basic properties of it. We study the closure, interior, base, continuity and compactness and properties of these concepts in fuzzy parametrized soft topological spaces.

In 1965, after Zadeh [32] generalized the usual notion of a set with the introduction of fuzzy set, the fuzzy set was carried out in the areas of general theories and applied to many real life problems in uncertain, ambiguous environment. In this manner, in 1968, Chang [13] gave the definition of fuzzy topology and introduced the many topological notions in fuzzy setting.

In 1999, Molodtsov [26] introduced the concept of soft set theory which is a completely new approach for modelling uncertainty and pointed out several directions for the applications of soft sets, such as game theory, perron integrations, smoothness of functions and so on. To improve this concept, many researchers applied this concept on topological spaces (e.g. [6, 15, 16, 19, 27, 33]), group theory, ring theory (e.g. [1, 2, 12, 17, 21]), and also decision making problems (e.g. [10, 11, 14, 23]).

Recently, researchers have combined fuzzy set and soft set to generalize the spaces and to solve more complicated problems. By this way, many interesting applications of soft set theory have been expanded. First combination of fuzzy set and soft set is fuzzy soft set and it was given by Maji and et al [24]. Then fuzzy soft set theory has been applied in several directions, such as topology (e.g. [3, 5, 29, 30]), various algebraic structures (e.g. [4, 20]) and especially decision making (e.g. [18, 22, 28, 31]). Another combination of fuzzy set and soft set was given by Çağman and et al. [7] and called it as fuzzy parametrized soft set (as shortly FP-soft set). In that paper, Çağman and et al. defined operations on FP-soft sets and improved several results. After that, Çağman and Deli [8, 9] applied FP-soft sets to define some decision making methods and applied these methods to problems that contain uncertainties and fuzzy object.

In the present paper, we consider the topological structure of FP-soft sets. Firstly, we give some basic ideas of FP-soft sets and also studied results. We define FP-soft quasi-coincidence, as a generalization of quasi-coincidence in fuzzy manner [25] and use this notion to characterize concepts of FP-soft closure and FP-soft base in FP-soft topological spaces. We also introduce the notion of mapping on FP-soft classes and investigate the properties of FP-soft images and FP-soft inverse images

2000 *Mathematics Subject Classification.* 03E72, 54C05, 54D30.

*Key words and phrases.* fuzzy parametrized soft set, fuzzy parametrized soft mapping, topology.

of FP-soft sets. We define FP-soft topology in Chang's sense. We study the FP-soft closure and FP-soft interior operators and properties of these concepts. Lastly we define FP-soft continuous mappings and we show that image of a FP-soft compact space is also FP-soft compact.

## 1. Preliminaries

Throughout this paper $X$ denotes initial universe, $E$ denotes the set of all possible parameters which are attributes, characteristic or properties of the objects in $X$, and the set of all subsets of $X$ will be denoted by $P(X)$.

**Definition 1.** [32] *A fuzzy set $A$ in $X$ is a function defined as follows:*
$A = \{\mu_A(x)/x : x \in X\}$
*where $\mu_A : X \to [0,1]$.*

*Here $\mu_A$ is called the membership function of $A$, and the value $\mu_A(x)$ is called the grade of membership of $x \in X$. This value represents the degree of $x$ belonging to the fuzzy set $A$.*

*A fuzzy point in $X$, whose value is $\alpha$ $(0 < \alpha \leq 1)$ at the support $x \in X$, is denoted by $x_\alpha$. A fuzzy point $x_\alpha \in A$, where $A$ is fuzzy set in $X$ iff $\alpha \leq \mu_A(x)$.*

**Definition 2.** [26] *A pair $(F, E)$ is called a soft set over $X$ if $F$ is a mapping defined by $F : E \to P(X)$.*

*In the other words, a soft set is a parametrized family of subsets of the set $X$. Each set $F(e)$, $e \in E$, from this family may be considered as the set of $e$-elements of the soft set $(F, E)$.*

**Definition 3.** [7] *Let $A$ be a fuzzy set over $E$. An FP-soft set $F_A$ on the universe $X$ is defined as follows:*
$F_A = \{(\mu_A(e)/e, f_A(e)) : e \in E, f_A(e) \in P(X), \mu_A(e) \in [0,1]\}$,
*where the function $f_A : E \to P(X)$ is called approximate function such that $f_A(e) = \varnothing$ if $\mu_A(e) = 0$, and the function $\mu_A : E \to [0,1]$ is called membership function of the set $A$.*

From now on, the set of all FP-soft sets over $X$ will be denoted by $FPS(X, E)$.

**Definition 4.** [7] *Let $F_A \in FPS(X, E)$.*

*(1) $F_A$ is called the empty FP-soft set if $\mu_A(e) = 0$ for all every $e \in E$, denoted by $F_\varnothing$.*

*(2) $F_A$ is called $A$-universal FP-soft set if $\mu_A(e) = 1$ and $f_A(e) = X$ for all $e \in A$, denoted by $F_{\widetilde{A}}$.*

*If $A = E$, then $A$-universal FP-soft set is called universal FP-soft set, denoted by $F_{\widetilde{E}}$.*

**Definition 5.** [7] *Let $F_A, F_B \in FPS(X, E)$.*

*(1) $F_A$ is called a FP-soft subset of $F_B$ if $A \leq B$ and $f_A(e) \subseteq f_B(e)$ for every $e \in E$ and we write $F_A \widetilde{\subset} F_B$.*

*(2) $F_A$ and $F_B$ are said to be equal, denoted by $F_A = F_B$ if $F_A \widetilde{\subset} F_B$ and $F_B \widetilde{\subset} F_A$.*

*(3) The union of $F_A$ and $F_B$, denoted by $F_A \widetilde{\cup} F_B$, is the FP-soft set, defined by the membership and approximate functions $\mu_{A\cup B}(e) = \max\{\mu_A(e), \mu_B(e)\}$ and $f_{A\cup B}(e) = f_A(e) \cup g_B(e)$ for every $e \in E$, respectively.*

*(4) The intersection of $F_A$ and $F_B$, denoted by $F_A \widetilde{\cap} G_B$, is the FP-soft set, defined by the membership and approximate functions $\mu_{A\cap B}(e) = \min\{\mu_A(e), \mu_B(e)\}$ and $f_{A\cap B}(e) = f_A(e) \cap g_B(e)$ for every $e \in E$, respectively.*

**Definition 6.** [7] *Let $F_A \in FPS(X,E)$. Then the complement of $F_A$, denoted by $F_A^c$, is the FP-soft set, defined by the membership and approximate functions $\mu_{A^c}(e) = 1 - \mu_A(e)$ and $f_A^c(e) = X - f_A(e)$ for every $e \in E$, respectively.*

*Clearly $(F_A^c)^c = F_A$, $F_{\widetilde{E}}^c = F_{\varnothing}$ and $F_{\varnothing}^c = F_{\widetilde{E}}$*

**Proposition 1.** [7] *Let $F_A$, $F_B$ and $F_C \in FPS(X,E)$. Then*

*(1) $(F_A \widetilde{\cup} F_B)^c = F_A^c \widetilde{\cap} F_B^c$*
*(2) $(F_A \widetilde{\cap} F_B)^c = F_A^c \widetilde{\cup} F_B^c$*
*(3) $F_A \widetilde{\cap} F_A = F_A$, $F_A \widetilde{\cup} F_A = F_A$*
*(4) $F_A \widetilde{\cap} F_{\varnothing} = F_{\varnothing}$, $F_A \widetilde{\cap} F_{\widetilde{E}} = F_A$*
*(5) $F_A \widetilde{\cap} F_B = F_B \widetilde{\cap} F_A$, $F_A \widetilde{\cup} F_B = F_B \widetilde{\cup} F_A$*
*(6) $F_A \widetilde{\cap}(F_B \widetilde{\cap} F_C) = (F_A \widetilde{\cap} F_B) \widetilde{\cap} F_C$, $F_A \widetilde{\cup}(F_B \widetilde{\cup} F_C) = (F_A \widetilde{\cup} F_B) \widetilde{\cup} F_C$*
*(7) $F_A \widetilde{\cup} F_{\varnothing} = F_A$, $F_A \widetilde{\cup} F_{\widetilde{E}} = F_{\widetilde{E}}$*

## 2. some properties of FP-soft sets and FP-soft mappings

**Definition 7.** *Let $J$ be an arbitrary index set and $F_{A_i} \in FPS(X,E)$ for all $i \in J$.*

*(1) The union of $F_{A_i}$'s, denoted by $\widetilde{\bigcup_{i\in J}} F_{A_i}$, is the FP-soft set, defined by the membership and approximate functions $\mu_{\bigcup_{i\in J} A_i}(e) = \sup_{i\in J}\{\mu_{A_i}(e)\}$ and $f_{\bigcup_{i\in J} A_i}(e) = \bigcup_{i\in J} f_{A_i}(e)$ for every $e \in E$, respectively.*

*(2) The intersection of $F_{A_i}$'s, denoted by $\widetilde{\bigcap_{i\in J}} F_{A_i}$, is the FP-soft set, defined by the membership and approximate functions $\mu_{\bigcap_{i\in J} A_i}(e) = \inf_{i\in J}\{\mu_{A_i}(e)\}$ and $f_{\bigcap_{i\in J} A_i}(e) = \bigcap_{i\in J} f_{A_i}(e)$ for every $e \in E$, respectively.*

**Proposition 2.** *Let $J$ be an arbitrary index set and $F_{A_i} \in FPS(X,E)$ for all $i \in J$. Then*

*(1) $(\widetilde{\bigcup_{i\in J}} F_{A_i})^c = \widetilde{\bigcap_{i\in J}} F_{A_i}^c$.*
*(2) $(\widetilde{\bigcap_{i\in J}} F_{A_i})^c = \widetilde{\bigcup_{i\in J}} F_{A_i}^c$.*

*Proof.* (1) Put $F_B = (\widetilde{\bigcup_{i\in J}} F_{A_i})^c$ and $F_C = \widetilde{\bigcap_{i\in J}} F_{A_i}^c$. Then for all $e \in E$,

$$\mu_B(e) = 1 - \mu_{\bigcup_{i\in J} A_i}(e) = 1 - \sup_{i\in J}\{\mu_{A_i}(e)\} = \inf_{i\in J}\{1 - \mu_{A_i}(e)\} = \inf_{i\in J}\{\mu_{A_i^c}(e)\} = \mu_C(e)$$

and

$$f_B(e) = X - f_{\bigcup_{i\in J} A_i}(e) = X - \bigcup_{i\in J} f_{A_i}(e) = \bigcap_{i\in J}(X - f_{A_i}(e)) = \bigcap_{i\in J} f_{A_i^c}(e) = f_{\bigcap_{i\in J} A_i^c}(e) = f_C(e).$$

This completes the proof. The other can be proved similarly □

**Definition 8.** *The FP-soft set $F_A \in FPS(X,E)$ is called FP-soft point if $A$ is fuzzy singleton and $f_A(e) \in P(X)$ for $e \in suppA$. If $A = \{e\}$, $\mu_A(e) = \alpha \in (0,1]$, then we denote this FP-soft point by $e_\alpha^f$.*

**Definition 9.** *Let $e_\alpha^f$, $F_A \in FPS(X,E)$. We say that $e_\alpha^f \widetilde{\in} F_A$ read as $e_\alpha^f$ belongs to the FP-soft set $F_A$ if $\alpha \le \mu_A(e)$ and $f(e) \subseteq f_A(e)$.*

**Proposition 3.** *Every non empty FP-soft set $F_A$ can be expresssed as the union of all the FP-soft points which belong to $F_A$.*

*Proof.* This follows from the fact that any fuzzy set is the union of fuzzy points which belong to it [25]. □

**Definition 10.** *Let $F_A$, $F_B \in FPS(X,E)$. $F_A$ is said to be FP-soft quasi-coincident with $F_B$, denoted by $F_A q F_B$, if there exists $e \in E$ such that $\mu_A(e) + \mu_B(e) > 1$ or $f_A(e)$ is not subset of $f_B^c(e)$. If $F_A$ is not FP-soft quasi-coincident with $F_B$, then we write $F_A \overline{q} F_B$.*

**Definition 11.** *Let $e_\alpha^f$, $F_A \in FPS(X,E)$. $e_\alpha^f$ is said to be FP-soft quasi-coincident with $F_A$, denoted by $e_\alpha^f q F_A$, if $\alpha + \mu_A(e) > 1$ or $f(e)$ is not subset of $f_A^c(e)$. If $e_\alpha^f$ is not FP-soft quasi-coincident with $F_A$, then we write $e_\alpha^f \overline{q} F_A$.*

**Proposition 4.** *Let $F_A$, $F_B \in FPS(X,E)$, Then the following are true.*

*(1) $F_A \widetilde{\subseteq} F_B \Leftrightarrow F_A \overline{q} F_B^c$.*
*(2) $F_A q F_B \Rightarrow F_A \widetilde{\cap} F_B \neq F_\varnothing$.*
*(3) $F_A \overline{q} F_A^c$.*
*(4) $F_A q F_B \Leftrightarrow$ there exists an $e_\alpha^f \widetilde{\in} F_A$ such that $e_\alpha^f q F_B$.*
*(5) $e_\alpha^f \widetilde{\in} F_A^c \Leftrightarrow e_\alpha^f \overline{q} F_A$.*
*(6) $F_A \widetilde{\subseteq} F_B \Rightarrow$ If $e_\alpha^f q F_A$, then $e_\alpha^f q F_B$ for all $e_\alpha^f \in FPS(X,E)$.*

*Proof.* (1)

$$\begin{aligned} F_A \widetilde{\subseteq} F_B &\Leftrightarrow \text{for all } e \in E,\ \mu_A(e) \leq \mu_B(e) \text{ and } f_A(e) \subseteq f_B(e) \\ &\Leftrightarrow \text{for all } e \in E\ ,\ \mu_A(e) - \mu_B(e) \leq 0 \text{ and } f_A(e) \subseteq f_B(e) \\ &\Leftrightarrow \text{for all } e \in E,\ \mu_A(e) + 1 - \mu_B(e) \leq 1 \text{ and } f_A(e) \subseteq f_B(e) \\ &\Leftrightarrow F_A \overline{q} F_B^c \end{aligned}$$

(2) Let $F_A q F_B$. Then there exists an $e \in E$ such that $\mu_A(e) + \mu_B(e) > 1$ or $f_A(e)$ is not subset of $f_B^c(e)$. If $\mu_A(e) + \mu_B(e) > 1$, $A \wedge B \neq 0_E$ and the proof is easy. If $f_A(e)$ is not subset of $f_B^c(e)$, then $f_A(e) \cap f_B(e) \neq \varnothing$. Hence $F_A \widetilde{\cap} F_B \neq F_\varnothing$.

(3) Suppose that $F_A q F_A^c$. Then there exists $e \in E$ such that $\mu_A(e) + \mu_{A^c}(e) > 1$ or $f_A(e)$ is not subset $(f_A^c(e))^c$. But this is impossible.

(4) If $F_A q F_B$, then there exist an $e \in E$ such that $\mu_A(e) + \mu_B(e) > 1$ or $f_A(e)$ is not subset of $f_B^c(e)$. Put $\alpha = \mu_A(e)$ and $f(e) = f_A(e)$. Then we have $e_\alpha^f \widetilde{\in} F_A$ and $e_\alpha^f q F_B$.

Conversely, suppose that $e_\alpha^f q F_B$ for some $e_\alpha^f \widetilde{\in} F_A$. Then $\alpha + \mu_B(e) > 1$ or $f(e)$ is not subset of $f_B^c(e)$. Therefore, we have $\mu_A(e) + \mu_B(e) > 1$ or $f_A(e)$ is not subset of $f_B^c(e)$ for $e \in E$ . This shows that $F_A q F_B$.

(5) It is obvious from (1).

(6) Let $e_\alpha^f$, $F_A \in FPS(X,E)$ and $e_\alpha^f q F_A$. Then $\alpha + \mu_A(e) > 1$ or $f(e)$ is not subset of $f_A^c(e)$. Since $F_A \widetilde{\subseteq} F_B$, $\alpha + \mu_B(e) > 1$ or $f(e)$ is not subset of $f_B^c(e)$. Hence we have $e_\alpha^f q F_B$. □

**Proposition 5.** *Let $\{F_{A_i} : i \in J\}$ be a family of FP-soft sets in $FPS(X,E)$ where $J$ is an index set. Then $e_\alpha^f$ is FP-soft quasi-coincident with $\widetilde{\cup}_{i \in J} F_{A_i}$ if and only if there exists some $F_{A_i} \in \{F_{A_i} : i \in J\}$ such that $e_\alpha^f q F_{A_i}$.*

*Proof.* Obvious. □

**Definition 12.** *Let $FPS(X,E)$ and $FPS(Y,K)$ be families of all FP-soft sets over $X$ and $Y$, respectively. Let $u : X \to Y$ and $p : E \to K$ be two functions. Then a FP-soft mapping $f_{up} : FPS(X,E) \to FPS(Y,K)$ is defined as:*

*(1) for $F_A \in FPS(X,E)$, the image of $F_A$ under the FP-soft mapping $f_{up}$ is the FP-soft set $G_S$ over $Y$ defined by the approximate function, $\forall k \in K$,*

$$g_S(k) = \begin{cases} \bigcup\limits_{e \in p^{-1}(k)} u(f_A(e)), & \text{if } p^{-1}(k) \neq \varnothing; \\ \varnothing, & \text{otherwise.} \end{cases}$$

*where $p(A) = S$ is fuzzy set in $K$.*

*(2) for $G_S \in FPS(Y,K)$, then the pre-image of $G_S$ under the FP-soft mapping $f_{up}$ is the FP-soft set $F_A$ over $X$ defined by the approximate function, $\forall e \in E$*

*$f_A(e) = u^{-1}(g_S(p(e)))$ where $p^{-1}(S) = A$ is fuzzy set in $E$.*

If $u$ and $p$ is injective, then the FP-soft mapping $f_{up}$ is said to be injective. If $u$ and $p$ is surjective, then the FP-soft mapping $f_{up}$ is said to be surjective. The FP-soft mapping $f_{up}$ is called constant, if $u$ and $p$ are constant.

**Theorem 1.** *Let $X$ and $Y$ crips sets $F_A$, $F_{A_i} \in FPS(X,E)$, $G_S$, $G_{S_i} \in FPS(Y,K)$ $\forall i \in J$, where $J$ is an index set. Let $f_{up} : FPS(X,E) \to FPS(Y,K)$ be a FP-soft mapping. Then,*

*(1) If $F_{A_1} \widetilde{\subset} F_{A_2}$ then $f_{up}(F_{A_1}) \widetilde{\subset} f_{up}(F_{A_2})$.*
*(2) If $G_{S_1} \widetilde{\subset} G_{S_2}$ then $f_{up}^{-1}(G_{S_1}) \widetilde{\subset} f_{up}^{-1}(G_{S_2})$.*
*(3) $F_A \widetilde{\subset} f_{up}^{-1}(f_{up}(F_A))$, the equality holds if $f_{up}$ is injective.*
*(4) $f_{up}(f_{up}^{-1}(G_S)) \widetilde{\subset} G_S$, the equality holds if $f_{up}$ is surjective.*
*(5) $f_{up}(\widetilde{\cup}_{i\in J} F_{A_i}) = \widetilde{\cup}_{i\in J} f_{up}(F_{A_i})$.*
*(6) $f_{up}(\widetilde{\cap}_{i\in J} F_{A_i}) \widetilde{\subset} \widetilde{\cap}_{i\in J} f_{up}(F_{A_i})$, the equality holds if $f_{up}$ is injective.*
*(7) $f_{up}^{-1}(\widetilde{\cup}_{i\in J} G_{S_i}) = \widetilde{\cup}_{i\in J} f_{up}^{-1}(G_{S_i})$.*
*(8) $f_{up}^{-1}(\widetilde{\cap}_{i\in J} G_{S_i}) = \widetilde{\cap}_{i\in J} f_{up}^{-1}(G_{S_i})$.*
*(9) $(f_{up}^{-1}(G_S))^c = f_{up}^{-1}(G_S^c)$.*
*(10) $(f_{up}(F_A))^c \widetilde{\subset} f_{up}(F_A^c)$.*
*(11) $f_{up}^{-1}(G_{\widetilde{K}}) = F_{\widetilde{E}}$.*
*(12) $f_{up}^{-1}(G_{\varnothing}) = F_{\varnothing}$.*
*(13) $f_{up}(F_{\widetilde{E}}) \widetilde{\subset} G_{\widetilde{K}}$, the equality holds if $f_{up}$ is surjective.*
*(14) $f_{up}(F_{\varnothing}) = G_{\varnothing}$.*

*Proof.* We only prove (3),(5),(7),(9),(11) and (12). The others can be proved similarly.

(3) Put $G_S = f_{up}(F_A)$ and $F_B = f_{up}^{-1}(G_S)$. Since $A \leq p^{-1}(p(A)) = p^{-1}(S) = B$, It is sufficient to show $f_A(e) \subseteq f_B(e)$ for all $e \in E$,

$$\begin{aligned} f_B(e) &= u^{-1}(g_S(p(e))) \\ &= u^{-1}(\cup_{e\in p^{-1}(p(e))} u(f_A(e))) \\ &= \cup_{e\in p^{-1}(p(e))} u^{-1}(u(f_A(e))) \\ &\supseteq f_A(e) \end{aligned}$$

This completes the proof.

(5) Put $G_{S_i} = f_{up}(F_{A_i})$ and $G_S = f_{up}(\widetilde{\cup}_{i\in J}(F_{A_i}))$. Then $S = p(\vee A_i) = \vee p(A_i) = \vee S_i$ and for all $k \in K$,

$$\begin{aligned} g_S(k) &= \begin{cases} \cup_{e\in p^{-1}(k)} u(\underset{i\in J}{\cup} f_{A_i}(e)) & \text{if } p^{-1}(k) \neq \varnothing; \\ \varnothing & \text{otherwise.} \end{cases} \\ &= \begin{cases} \cup_{e\in p^{-1}(k)} \underset{i\in J}{\cup} u(f_{A_i}(e)) & \text{if } p^{-1}(k) \neq \varnothing; \\ \varnothing & \text{otherwise.} \end{cases} \\ &= \begin{cases} \underset{i\in J}{\cup} \cup_{e\in p^{-1}(k)} u(f_{A_i}(e)) & \text{if } p^{-1}(k) \neq \varnothing; \\ \varnothing & \text{otherwise.} \end{cases} \\ &= \underset{i\in J}{\cup} \begin{cases} \cup_{e\in p^{-1}(k)} u(f_{A_i}(e)) & \text{if } p^{-1}(k) \neq \varnothing; \\ \varnothing & \text{otherwise.} \end{cases} \\ &= \cup_{i\in J} g_{S_i}(k) \end{aligned}$$

This completes the proof.

(7) Put $F_{A_i} = f_{up}^{-1}(G_{S_i})$ and $F_A = f_{up}^{-1}(\widetilde{\cup}_{i\in J} G_{S_i})$. Then $A = p^{-1}(\vee S_i) = \vee p^{-1}(S_i) = \vee A_i$ and for all $e \in E$,

$$\begin{aligned} f_A(e) &= u^{-1}(\underset{i\in I}{\cup} g_{S_i}(p(e))) \\ &= \underset{i\in I}{\cup} u^{-1}(g_{S_i}(p(e))) \\ &= \cup_{i\in J} f_{A_i}(e) \end{aligned}$$

This completes the proof.

(9) Put $f_{up}^{-1}(G_S) = F_A$ and $f_{up}^{-1}(G_S^c) = F_B$. Then for all $e \in E$, $f_B(e) = f_{p^{-1}(S^c)}(e) = f_{(p^{-1}(S))^c}(e) = f_{A^c}(e)$ where $p^{-1}(S)$ and $p^{-1}(S^c)$ are fuzzy sets over $E$. This shows that the approximate functions of $F_B$ and $F_A^c$ are equal. This completes the proof.

(11) Put $F_A = f_{up}^{-1}(G_{\widetilde{K}})$. Then for all $e \in E$, $f_A(e) = u^{-1}(G_{\widetilde{K}}(p(e))) = u^{-1}(Y) = X = f_E(e)$. This shows that $F_A = F_{\widetilde{E}}$.

(12) Since $p^{-1}(K)$ is fuzzy empty set i.e. $0_E$, the proof is clear. □

## 3. FP-soft topological spaces

**Definition 13.** *A FP-soft topological space is a pair $(X, \tau)$ where $X$ is a nonempty set and $\tau$ is a family of FP-soft sets over $X$ satisfying the following properties:*

*(T1) $F_\varnothing, F_{\widetilde{E}} \in \tau$*

*(T2) If $F_A$, $F_B \in \tau$ , then $F_A \widetilde{\cap} F_B \in \tau$*

*(T3) If $F_{A_i} \in \tau$ ,$\forall i \in J$, then $\widetilde{\cup}_{i\in J} F_{A_i} \in \tau$.*

*$\tau$ is called a topology of FP-soft sets on $X$. Every member of $\tau$ is called FP-soft open in $(X, \tau)$. $F_B$ is called FP-soft closed in $(X, \tau)$ if $F_B^c \in \tau$.*

**Example 1.** *$\tau_{indiscrete} = \{F_\varnothing, F_{\widetilde{E}}\}$ is a FP-soft topology on $X$.*

*$\tau_{discrete} = FPS(X, E)$ is a FP-soft topology on $X$.*

**Example 2.** *Assume that $X = \{x_1, x_2, x_3, x_4\}$ is a universal set and $E = \{e_1, e_2, e_3\}$ is a set of parameters. If*

$F_{A_1} = \{((e_1)_{0,2}, \{x_1, x_3\}), ((e_2)_{0,3}, \{x_1, x_4\}), ((e_3)_{0,4}, \{x_2\})\}$

$F_{A_2} = \{((e_1)_{0,2}, \{x_1, x_2, x_3\}), ((e_2)_{0,5}, \{x_1, x_4\}), ((e_3)_{0,4}, \{x_1, x_2\})\}$

$F_{A_3} = \{((e_1)_{0,7}, \{x_1, x_3\}), ((e_2)_{0,3}, X), ((e_3)_{0,9}, \{x_2, x_3\})\}$

$F_{A_4} = \{((e_1)_{0,7}, \{x_1, x_2, x_3\}), ((e_2)_{0,5}, X), ((e_3)_{0,9}, \{x_2, x_3\})\}$

*then $\tau = \{F_\varnothing, F_{A_1}, F_{A_2}, F_{A_3}, F_{A_4}, F_{\widetilde{E}}\}$ is a FP-soft topology on $X$.*

**Theorem 2.** *Let $(X, \tau)$ a FP-soft topological space and $\tau'$ denote family of all closed sets. Then;*

*(1) $F_\varnothing, F_{\widetilde{E}} \in \tau'$*

*(2) If $F_A$, $F_B \in \tau'$, then $F_A \widetilde{\cup} F_B \in \tau'$*

*(3) If $F_{A_i} \in \tau'$, $\forall i \in J$, then $\widetilde{\cap}_{i\in J} F_{A_i} \in \tau'$.*

*Proof.* Straightforward. □

**Definition 14.** *Let $(X, \tau)$ be a FP-soft topological space and $F_A \in FPS(X, E)$. The FP-soft closure of $F_A$ in $(X, \tau)$, denoted by $\overline{F_A}$, is the intersection of all FP-soft closed supersets of $F_A$.*

*Clearly, $\overline{F_A}$ is the smallest FP-soft closed set over $X$ which contains $F_A$, and $\overline{F_A}$ is closed.*

**Theorem 3.** *Let $(X, \tau)$ be a FP-soft topological space and $F_A$, $F_B \in FPS(X, E)$. Then,*

*(1)* $\overline{F_{\varnothing}} = F_{\varnothing}$ *and* $\overline{F_{\widetilde{E}}} = F_{\widetilde{E}}$.
*(2)* $F_A \widetilde{\subset} \overline{F_A}$.
*(3)* $\overline{\overline{F_A}} = \overline{F_A}$.
*(4) If* $F_A \widetilde{\subset} F_B$, *then* $\overline{F_A} \widetilde{\subset} \overline{F_B}$.
*(5)* $F_A$ *is a FP-soft closed set if and only if* $F_A = \overline{F_A}$.
*(6)* $\overline{F_A \widetilde{\cup} F_B} = \overline{F_A} \widetilde{\cup} \overline{F_B}$.

*Proof.* (1),(2),(3) and (4) are obvious from the definition of FP-soft closure.

(5) Let $F_A$ be a FP-soft closed set. Since $\overline{F_A}$ is the smallest FP-soft closed set which contains $F_A$, then $\overline{F_A} \widetilde{\subset} F_A$. Therefore, $F_A = \overline{F_A}$.

(6) Since $F_A \widetilde{\subset} F_A \widetilde{\cup} F_B$ and $F_B \widetilde{\subset} F_A \widetilde{\cup} F_B$, then, by (4), $\overline{F_A} \widetilde{\subset} \overline{F_A \widetilde{\cup} F_B}$ , $\overline{F_B} \widetilde{\subset} \overline{F_A \widetilde{\cup} F_B}$ and hence $\overline{F_A} \widetilde{\cup} \overline{F_B} \widetilde{\subset} \overline{F_A \widetilde{\cup} F_B}$.

Conversely, since $\overline{F_A}$, $\overline{F_B}$ are FP-soft closed sets, $\overline{F_A} \widetilde{\cup} \overline{F_B}$ is a FP-soft closed set. Again since $F_A \widetilde{\cup} F_B \widetilde{\subset} \overline{F_A} \widetilde{\cup} \overline{F_B}$, by (4), then $\overline{F_A \widetilde{\cup} F_B} \widetilde{\subset} \overline{F_A} \widetilde{\cup} \overline{F_B}$. □

**Definition 15.** *Let* $(X, \tau)$ *be a FP-soft topological space. A FP-soft set* $F_A$ *in* $FPS(X, E)$ *is called FP-Q-neighborhood (briefly, FP-Q-nbd) of a FP-soft set* $F_B$ *if there exists a FP-soft open set* $F_C$ *in* $\tau$ *such that* $F_B q F_C$ *and* $F_C \widetilde{\subseteq} F_A$.

**Theorem 4.** *Let* $e^f_\alpha$, $F_A \in FPS(X, E)$. *Then* $e^f_\alpha \widetilde{\in} \overline{F_A}$ *if and only if each FP-Q-nbd of* $e^f_\alpha$ *is FP-soft quasi-coincident with* $F_A$.

*Proof.* Let $e^f_\alpha \widetilde{\in} \overline{F_A}$. Suppose that $F_C$ is a FP-Q-nbd of $e^f_\alpha$ and $F_C \overline{q} F_A$. Then there exists a FP-soft open set $F_B$ such that $e^f_\alpha q F_B \widetilde{\subseteq} F_C$. Since $F_C \overline{q} F_A$, by Proposition 4(1), $F_A \widetilde{\subseteq} F^c_C \widetilde{\subseteq} F^c_B$. Again since $e^f_\alpha q F_B$, $e^f_\alpha$ does not belong to $F^c_B$. This is a contradiction with $\overline{F_A} \widetilde{\subseteq} F^c_B$.

Conversely, Let each Q-nbd of $e^f_\alpha$ be FP-soft quasi-coincident with $F_A$. Suppose that $e^f_\alpha$ does not belong to $\overline{F_A}$. Then there exists a FP-soft closed set $F_B$ which is containing $F_A$ such that $e^f_\alpha$ does not belong to $F_B$. By Proposition 4(5), we have $e^f_\alpha q F^c_B$. Then $F^c_B$ is a FP-Q-nbd of $e^f_\alpha$ and by Proposition 4(1), $F_A \overline{q} F^c_B$. This is a contradiction with the hypothesis. □

**Definition 16.** *Let* $(X, \tau)$ *be a FP-soft topological space and* $F_A \in FPS(X, E)$. *The FP-soft interior of* $F_A$ *denoted by* $F^\circ_A$ *is the union of all FP-soft open subsets of* $F_A$.

*Clearly,* $F^\circ_A$ *is the largest fuzzy soft open set contained in* $F_A$ *and* $F^\circ_A$ *is FP-soft open.*

**Theorem 5.** *Let* $(X, \tau)$ *be a FP-soft topological space and* $F_A$, $F_B \in FPS(X, E)$. *Then,*
*(1)* $(F_{\varnothing})^\circ = F_{\varnothing}$ *and* $(F_{\widetilde{E}})^\circ = F_{\widetilde{E}}$.
*(2)* $F^\circ_A \widetilde{\subset} F_A$.
*(3)* $(F^\circ_A)^\circ = F^\circ_A$.
*(4) If* $F_A \widetilde{\subset} F_B$, *then* $F^\circ_A \widetilde{\subset} F^\circ_B$.
*(5)* $F_A$ *is a FP-soft open set if and only if* $F_A = F^\circ_A$.
*(6)* $\left(F_A \widetilde{\cap} F_B\right)^\circ = F^\circ_A \widetilde{\cap} F^\circ_B$.

*Proof.* Similar to that of Theorem 3. □

**Theorem 6.** *Let* $(X, \tau)$ *be a FP-soft topological space and* $F_A \in FPS(X, E)$. *Then,*
*(1)* $\left(F^\circ_A\right)^c = \overline{F^c_A}$.
*(2)* $\left(\overline{F_A}\right)^c = (F^c_A)^\circ$.

*Proof.* We only prove (1). The other is similar.

$$\begin{aligned}(F_A^\circ)^c &= (\widetilde{\cup}\{F_B \,|F_B \in \tau,\ F_A\widetilde{\subset}F_B\ \})^c\\ &= \widetilde{\cap}\{F_B^c \,|F_B \in \tau,\ F_A\widetilde{\subset}F_B\ \}\\ &= \widetilde{\cap}\{F_B^c \,|F_B^c \in \tau',\ F_B^c\widetilde{\subset}F_A^c\ \}\\ &= \overline{F_A^c}\end{aligned}$$

□

**Theorem 7.** *Let $c : FPS(X,E) \to FPS(X,E)$ be an operator satisfying the following:*

*(c1) $c(F_\varnothing) = F_\varnothing$.*

*(c2) $F_A\widetilde{\subseteq}c(F_A), \forall F_A \in FPS(X,E)$.*

*(c3) $c(F_A\widetilde{\cup}F_B) = c(F_A)\widetilde{\cup}c(F_B), \forall F_A, F_B \in FPS(X,E)$.*

*(c4) $c(c(F_A)) = c(F_A), \forall F_A \in FPS(X,E)$.*

*Then we can associate FP-soft topology in the following way:*

$\tau = \{F_A^c \in FPS(X,E) | c(F_A) = F_A\}$.

*Moreover with this FP-soft topology $\tau$, $\overline{F_A} = c(F_A)$ for every $F_A \in FPS(X,E)$.*

*Proof.* (T1) By (c1), $F_\varnothing^c = F_{\widetilde{E}} \in \tau$. By (c2) $F_{\widetilde{E}}\widetilde{\subseteq}c(F_{\widetilde{E}})$, so $c(F_{\widetilde{E}}) = F_{\widetilde{E}}$ and $F_\varnothing \in \tau$.

(T2) Let $F_A$, $F_B \in \tau$. By the definition of $\tau$, $c(F_A^c) = F_A^c$ and $c(F_B^c) = F_B^c$. By (c3), $c((F_A\widetilde{\cap}F_B)^c) = c(F_A^c\widetilde{\cup}F_B^c) = c(F_A^c)\widetilde{\cup}c(F_B^c) = F_A^c\widetilde{\cup}F_B^c = (F_A\widetilde{\cap}F_B)^c$. So $F_A\widetilde{\cap}F_B \in \tau$.

(T3) Let $\{F_{A_i} | i \in J\} \subset \tau$. Since $c$ is order preserving and $\widetilde{\bigcap_{i\in J}} F_{A_i}^c \widetilde{\subseteq} F_{A_k}^c, \forall k \in J$, then $c(\widetilde{\bigcap_{i\in J}} F_{A_i}^c)\widetilde{\subseteq}c(F_{A_k}^c) = F_{A_k}^c$. Then we have $c(\widetilde{\bigcap_{i\in J}} F_{A_i}^c)\widetilde{\subseteq}\widetilde{\bigcap_{i\in J}} F_{A_i}^c$. Conversely, by (c2) we have $\widetilde{\bigcap_{i\in J}} F_{A_i}^c \widetilde{\subseteq} c(\widetilde{\bigcap_{i\in J}} F_{A_i}^c)$. Hence, $c((\widetilde{\bigcup_{i\in J}} F_{A_i})^c) = c(\widetilde{\bigcap_{i\in J}} F_{A_i}^c) = \widetilde{\bigcap_{i\in J}} F_{A_i}^c = (\widetilde{\bigcup_{i\in J}} F_{A_i})^c$ and $\widetilde{\bigcup_{i\in J}} F_{A_i} \in \tau$.

Now we will show that with this FP-soft topology $\tau$, $\overline{F_A} = c(F_A)$ for every $F_A \in FPS(X,E)$. Let $F_A \in FPS(X,E)$. Since $(\overline{F_A})^c \in \tau$, then $c(\overline{F_A}) = \overline{F_A}$. Since $c$ is order preserving $c(F_A)\widetilde{\subseteq}c(\overline{F_A}) = \overline{F_A}$. Conversely, by (c4) we have $(c(F_A))^c \in \tau$. Then since $F_A\widetilde{\subseteq}c(F_A)$ and $\overline{F_A}$ is the smallest FP-soft closed set over $X$ which contains $F_A$, $\overline{F_A}\widetilde{\subseteq}c(F_A)$. □

The operator $c$ is called the FP-soft closure operator.

**Remark 1.** *By Theorem 3(1),(2),(3) and (6) and Theorem 7, we see that with a FP-soft closure operator we can associate a FP-soft topology and conversely with a given FP-soft topology we can associte a FP-soft closure operator.*

**Theorem 8.** *Let $i : FPS(X,E) \to FPS(X,E)$ be an operator satisfying the following:*

*(i1) $i(F_{\widetilde{E}}) = F_{\widetilde{E}}$.*

*(i2) $i(F_A)\widetilde{\subseteq}F_A$, $\forall F_A \in FPS(X,E)$.*

*(i3) $i(F_A\widetilde{\cap}F_B) = i(F_A)\widetilde{\cap}i(F_B)$, $\forall F_A, F_B \in FPS(X,E)$.*

*(i4) $i(i(F_A)) = i(F_A)$, $\forall F_A \in FPS(X,E)$.*

*Then we can associate a FP-soft topology in the following way:*

$\tau = \{F_A \in FPS(X,E) | i(F_A) = F_A\}$.

*Moreover, with this fuzzy soft topology $\tau$,$(F_A)^\circ = i(F_A)$ for every $F_A \in FPS(X,E)$.*

*Proof.* Similar to that of Theorem 7. □

The operator $i$ is called the FP-soft interior operator.

**Remark 2.** *By Theorem 5(1),(2),(3) and (6) and Theorem 8, we see that with a FP-soft interior operator we can associate a FP-soft topology and conversely with a given FP-soft topology we can associate a FP-soft interior operator.*

**Definition 17.** *Let $(X,\tau)$ be a FP-soft topological space. A subcollection $\mathcal{B}$ of $\tau$ is called a base for $\tau$ if every member of $\tau$ can be expressed as a union of members of $\mathcal{B}$.*

**Example 3.** *If we consider the FP-soft topology $\tau$ in Example 2, then one easily see that the family $\mathcal{B}=\{F_{\varnothing}, F_{A_1}, F_{A_2}, F_{A_3}, F_{\widetilde{E}}\}$ is a basis for $\tau$.*

**Proposition 6.** *Let $(X,\tau)$ be a FP-soft topological space and $\mathcal{B}$ is subfamily of $\tau$. $\mathcal{B}$ is a base for $\tau$ if and only if for each $e_\alpha^f$ in $FPS(X,E)$ and for each FP-soft open Q-nbd $F_A$ of $e_\alpha^f$, there exists a $F_B\in\mathcal{B}$ such that $e_\alpha^f q F_B \widetilde{\subseteq} F_A$.*

*Proof.* Let $\mathcal{B}$ be a base for $\tau$, $e_\alpha^f \widetilde{\in} FPS(X,E)$ and $F_A$ be a FP-soft open Q-nbd of $e_\alpha^f$. Then there exists a subfamily $\mathcal{B}'$ of $\mathcal{B}$ such that $F_A=\widetilde{\cup}\{F_B|\, F_B\in\mathcal{B}'\}$. Suppose that $e_\alpha^f \overline{q} F_B$ for all $F_B\in\mathcal{B}'$. Then $\alpha+\mu_B(e)\le 1$ and $f(e)\subseteq f_B^c(e)$ for every $F_B\in\mathcal{B}'$. Therefore, we have $\alpha+\mu_A(e)\le 1$ and $f(e)$ is not subset of $f_A(e)$ since $\mu_A(e)=\sup\{\mu_B(e)|\, F_B\in\mathcal{B}'\}$ and $f_A(e)=\cup f_B(e)$. This is contradiction.

Conversely, If $\mathcal{B}$ is not a base for $\tau$, then there exists a $F_A\in\tau$ such that $F_C=\widetilde{\cup}\{F_B\in\mathcal{B}: F_B\widetilde{\subseteq}F_A\}\neq F_A$. Since $F_C\neq F_A$, there exists $e\in E$ such that $\mu_C(e)<\mu_A(e)$ or $f_C(e)\subset f_A(e)$. Put $\alpha=1-\mu_C(e)$ and $f(e)=f_C^c(e)$. Then in both case, we obtain that $e_\alpha^f q F_A$ and $e_\alpha^f \overline{q} F_C$. Therefore, we have $\mu_B(e)+\alpha\le \mu_C(e)+\alpha=1$ and $f(e)\cap f_B(e)\subseteq f(e)\cap f_C(e)=\varnothing$; that is, $e_\alpha^f\overline{q}F_B$ for all $F_B\in\mathcal{B}$ which contained in $F_A$. This is a contradiction. □

**Definition 18.** *Let $(X,\tau_1)$ and $(Y,\tau_2)$ be two FP-soft topological spaces. A FP-soft mapping $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ is called FP-soft continuous if $f_{up}^{-1}(G_S)\in\tau_1$, $\forall G_S\in\tau_2$.*

**Example 4.** *Assume that $X=\{x_1,x_2,x_3\}$, $Y=\{y_1,y_2,y_3\}$ are two universal sets, $E=\{e_1,e_2\}$, $K=\{k_1,k_2\}$ are two parameter sets and $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ is a FP-soft mapping, where $u(x_1)=y_2$, $u(x_2)=y_1$, $u(x_3)=y_3$ and $p(e_1)=k_2$, $p(e_2)=k_1$. If we take $F_A=\{((e_1)_{0,3},\{x_2,x_3\}),((e_2)_{0,2},\{x_1,x_2\})\}$, $G_S=\{((k_1)_{0,2},\{y_1,y_2\}),((k_2)_{0,3},\{y_1,y_3\})\}$, $\tau_1=\{F_{\varnothing},F_{\widetilde{E}},F_A\}$ and $\tau_2=\{\widetilde{0}_K,\widetilde{1}_K,G_S\}$, then $f_{up}$ is a FP-soft continuous mapping.*

The constant mapping $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ not continuous in general.

**Example 5.** *Assume that $X=\{x_1,x_2,x_3\}$, $Y=\{y_1,y_2\}$ are two universal sets, $E=\{e_1,e_2\}$, $K=\{k_1,k_2\}$ are two parameter sets and $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ is a constant FP-soft mapping, where $u(x_1)=u(x_2)=u(x_3)=y_2$ and $p(e_1)=p(e_2)=k_1$. If we take $G_S=\{((k_1)_{0,2},\{y_1,y_2\}),((k_2)_{0,5},\{y_2,y_3\})\}$, $\tau_1=\{F_{\varnothing},F_{\widetilde{E}}\}$ and $\tau_2=\{F_{\varnothing},F_{\widetilde{K}},G_S\}$, $f_{up}$ is not a FP-soft continuous since $f_{up}^{-1}(G_S)\notin\tau_1$.*

Let $\alpha\in[0,1]$. We denote by $\alpha_E$ the constant fuzzy set on $E$, i.e $\mu_{\alpha_E}(e)=\alpha$ for all $e\in E$ and $\alpha\in[0,1]$.

**Definition 19.** *Let $F_A\in FPS(X,E)$. $F_A$ is called $\alpha-A$-universal FP-soft set if $\mu_A(e)=\alpha$ and $f_A(e)=X$ for all $e\in A$, denoted by $F_{\widetilde{\alpha}_A}$.*

**Definition 20.** *(see [5])A FP-soft topology is called enriched if it satisfies $F_{\widetilde{\alpha}_A}\in\tau$ and $F_{\widetilde{\alpha}_A}^c\in\tau$ for all $\alpha\in(0,1]$.*

**Theorem 9.** *Let $(X,\tau_1)$ be a enriched FP-soft topological space, $(Y,\tau_2)$ be a FP-soft topological space and $f_{up}:FPS(X,E)\to FPS(Y,K)$ be a constant FP-soft mapping. Then $f_{up}$ is FP-soft continuous.*

*Proof.* Let $G_S\in\tau_2$. Put $f_{up}^{-1}(G_S)=F_A$. Then $A=p^{-1}(S)=\alpha_E$ where $\alpha=\sup_{k\in K}\{\mu_S(k)\}$ and
$f_A(e)=u^{-1}(g_S(p(e))=\begin{cases} X & , g_S(p(e))\neq\varnothing \\ \varnothing & , \text{otherwise}\end{cases}$ for all $e\in E$. Hence $F_A=F_{\widetilde{\alpha}_E}\in\tau_1$ or $F_A=F^c_{\widetilde{\alpha}_E}\in\tau_1$ and so $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ is FP-soft continuous. □

**Theorem 10.** *Let $(X,\tau_1)$ and $(Y,\tau_2)$ be two FP-soft topological spaces and $f_{up}:FPS(X,E)\to FPS(Y,K)$ be a FP-soft mapping. Then the following are equivalent:*

*(1) $f_{up}$ is FP-soft continuous;*
*(2) $f_{up}^{-1}(G_S)$ is FP-soft closed for every FP- closed set $G_S$ over $Y$;*
*(3) $\overline{f_{up}(F_A)}\widetilde{\subset}f_{up}(\overline{F_A})$, $\forall F_A\in FPS(X,E)$;*
*(4) $\overline{f_{up}^{-1}(G_S)}\widetilde{\subset}f_{up}^{-1}(\overline{G_S})$, $\forall G_S\in FPS(Y,K)$;*
*(5) $f_{up}^{-1}(G_S^\circ)\widetilde{\subset}(f_{up}^{-1}(G_S))^\circ$, $\forall G_S\in FPS(Y,K)$.*

*Proof.* (1)⇒(2) It is obvious from Theorem 1 (9).

(2)⇒(3) Let $F_A\in FPS(X,E)$. Since $F_A\widetilde{\subset}f_{up}^{-1}(f_{up}(F_A))F_A\widetilde{\subset}f_{up}^{-1}(\overline{f_{up}(F_A)})\in\tau_1'$. Therefore we have $\overline{F_A}\widetilde{\subset}f_{up}^{-1}(\overline{f_{up}(F_A)})$. By Theorem 1 (4), we get $f_{up}(\overline{F_A})\widetilde{\subset}f_{up}(f_{up}^{-1}(\overline{f_{up}(F_A)}))\widetilde{\subset}\overline{f_{up}(F_A)}$.

(3)⇒(4) Let $G_S\in FPS(Y,K)$. If we choose $f_{up}^{-1}(G_S)$ instead of $F_A$ in (3), then $f_{up}(\overline{f_{up}^{-1}(G_S)})\widetilde{\subset}\overline{f_{up}(f_{up}^{-1}(G_S))}\widetilde{\subset}\overline{G_S}$. Hence by Theorem 1(3), $\overline{f_{up}^{-1}(G_S)}\widetilde{\subset}f_{up}^{-1}(f_{up}(\overline{f_{up}^{-1}(G_S)}))\widetilde{\subset}f_{up}^{-1}(\overline{G_S})$.

(4)⇔(5) These follow from Theorem 1 (9) and Theorem 6.

(5)⇒(1) Let $G_S\in\tau_2$. Since $G_S$ is a FP-soft open set, then $f_{up}^{-1}(G_S)=f_{up}^{-1}(G_S^\circ)\widetilde{\subset}(f_{up}^{-1}(G_S))^\circ\widetilde{\subset}f_{up}^{-1}(G_S)$. Consequently, $f_{up}^{-1}(G_S)$ is a FP-soft open and so $f_{up}$ is FP-soft continuous. □

**Theorem 11.** *Let $f_{up}:(X,\tau_1)\to(Y,\tau_2)$ be a FP-soft mapping and $\mathcal{B}$ be a base for $\tau_2$. Then $f_{up}$ is FP-soft continuous if and only if $f_{up}^{-1}(G_S)\in\tau_1$, $\forall G_S\in\mathcal{B}$.*

*Proof.* Straightforward. □

**Definition 21.** *A family $\mathcal{C}$ of FP-soft sets is a cover of a FP-soft set $F_A$ if $F_A\widetilde{\subseteq}\widetilde{\cup}\{F_{A_i}:F_{A_i}\in\mathcal{C},i\in J\}$. It is a FP-soft open cover if each member of $\mathcal{C}$ is a FP-soft open set. A subcover of $\mathcal{C}$ is a subfamily of $\mathcal{C}$ which is also a cover.*

**Definition 22.** *A family $\mathcal{C}$ of FP-soft sets has the finite intersection property if the intersection of the members of each finite subfamily of $\mathcal{C}$ is not empty FP-soft set.*

**Definition 23.** *A FP-soft topological space $(X,\tau)$ is FP-compact if each FP-soft open cover of $F_{\widetilde{E}}$ has a finite subcover.,*

**Example 6.** *Let $X=\{x_1,x_2,...\}$, $E=\{e_1,e_2,...\}$ and $F_{A_n}=\{((e_i)_{\frac{1}{n}},X-\{x_1,x_2,...x_n\}):i=1,2,...\}$. Then $\tau=\{F_{A_n}:n=1,2,...\}\cup\{F_\varnothing,F_{\widetilde{E}}\}$ is a FP-soft topology on $X$ and $(X,\tau)$ is FP-compact.*

**Theorem 12.** *A FP-soft topological space is FP-soft compact if and only if each family of FP-soft closed sets with the finite intersection property has a non empty FP-soft intersection.*

*Proof.* If $\mathcal{C}$ is a family of FP-soft sets in a FP-soft topological space $(X,\tau)$, then $\mathcal{C}$ is a cover of $F_{\widetilde{E}}$ if and only if one of the following conditions holds:

(1) $\widetilde{\cup}\{F_{A_i} : F_{A_i} \in \mathcal{C}, i \in J\} = F_{\widetilde{E}}$

(2) $(\widetilde{\cup}\{F_{A_i} : F_{A_i} \in \mathcal{C}, i \in J\})^c = F^c_{\widetilde{E}} = F_{\varnothing}$

(3) $\widetilde{\cap}\{F^c_{A_i} : F_{A_i} \in \mathcal{C}, i \in J\} = F_{\varnothing}$

Hence the FP-soft topological space is FP-soft compact if and only if each family of FP-soft open sets over $X$ such that no finite subfamily covers $F_{\widetilde{E}}$, fails to be a cover, and this is true if and only if each family of FP-soft closed sets which has the finite intersection property has a nonempty FP-soft intersection. □

**Theorem 13.** *Let $(X,\tau_1)$ and $(Y,\tau_2)$ be FP-soft topological spaces and $f_{up} : FPS(X,E) \to FPS(Y,K)$ be a FP-soft mapping. If $(X,\tau_1)$ is FP-soft compact and $f_{up}$ is FP-soft continuous surjection, then $(Y,\tau_2)$ is FP-soft compact.*

*Proof.* Let $\mathcal{C} = \{G_{S_i} : i \in J\}$ be a cover of $G_{\widetilde{K}}$ by FP-soft open sets. Then since $f_{up}$ is FP-soft continuous, the family of all FP-soft sets of the form $f^{-1}_{up}(G_S)$, for $G_S \in \mathcal{C}$, is a FP-soft open cover of $F_{\widetilde{E}}$ which has a finite subcover. However, since $f_{up}$ is surjective, then $f_{up}(f^{-1}_{up}(G_S) = G_S$ for any FP-soft set $G_S$ over $Y$. Thus, the family of images of members of the subcover is a finite subfamily of $\mathcal{C}$ which covers $G_{\widetilde{K}}$. Consequently, $(Y,\tau_2)$ is FP-soft compact. □

**Conclusion.** Topology is a branch of mathematics, whose concepts exist not only in almost all branches of mathematics, but also in many real life applications. In this paper, we introduce the topological structure of fuzzy parametrized soft sets and fuzzy parametrized soft mappings. We study some fundemental concepts in fuzzy parametrized soft topological spaces such as closures, interiors, bases, compactness and continuity. Some basic properties of these concepts are also presented. This paper will form the basis for further applications of topology.

## References


[1] Acar, U., Koyuncu, F. and Tanay, B. *Soft sets and soft rings*, Comput. Math. Appl. **59**, 3458-3463, 2010.

[2] Aktaş, H. and Çağman, N. *Soft sets and soft groups*, Inform. Sci. **177** (13), 2726-2735, 2007.

[3] Atmaca, S. and Zorlutuna, İ. *On fuzzy soft topological spaces*, Ann. Fuzzy Math. Inform. **5** (2), 377-386, 2013.

[4] Aygünoğlu, A. and Aygün, H. *Introduction to fuzzy soft groups*, Comput. Math. Appl. **58,** 1279-1286, 2009.

[5] Varol, B. P. and Aygün, H. *Fuzzy soft topology*, Hacet. J. Math. Stat. **41** (3), 407-419, 2012.

[6] Çağman, N., Karataş, S. and Enginoglu, S. *Soft topology*, Comput. Math. Appl. **62,** 351-358, 2011.

[7] Çağman, N., Çıtak, F. and Enginoglu, S. *FP-soft set theory and its applications*, Ann. Fuzzy Math. Inform. **2** (2), 219-226, 2011.

[8] Çağman, N. and Deli, İ. *Products of FP-soft sets and their applications*, Hacet. J. Math. Stat. **41** (3), 365-374, 2012.

[9] Çağman, N. and Deli, İ. Means *of FP-soft sets and their applications*, Hacet. J. Math. Stat. **41** (5), 615-625, 2012.

[10] Çağman, N. and Enginoglu, S. *Soft set theory and uni-int decision making*, Eur. J. Oper. Res. **207**, 848-855, 2010.

[11] Çağman, N. and Enginoglu, S. *Soft matrix theory and its decision making*, Comput. Math. Appl. **59**, 3308-3314, 2010.
[12] Çelik, Y., Ekiz, C. and Yamak, S. *A new view on soft rings*, Hacet. J. Math. Stat. **40** (2), 273-286, 2011.
[13] Chang, C. L. *Fuzzy topological spaces*, J. Math. Anal. Appl. **24**, 182-190, 1968.
[14] Chen, D., Tsang, E. C. C., Yeung, D. S. and Wang, X. *The parametrization reduction of soft sets and its applications*, Comput. Math. Appl. **49**, 757-763, 2005.
[15] B. Chen, *Soft semi-open sets and related properties in soft topological spaces*, Appl. Math. Inf. Sci., 7 (1) (2013), 287-294.
[16] B. Chen, *Some local properties of soft semi-open sets,* Discrete Dyn. Nat. Soc., Article ID 298032, 6 pages, 2013.
[17] Feng, F., Jun, Y.B. and Zhao, X. *Soft semirings*, Comput. Math. Appl. **56,** 2621-2628, 2008.
[18] Feng, F., Jun, Y.B., Liu, X. and Li, L.F. *An adjustable approach to fuzzy soft set based decision making*, J. Comput. Appl. Math. **234**, 10-20, 2010.
[19] Georgiou, D. N., Megaritis, A.C. and Petropoulos, V.I. *On soft topological spaces*, Appl. Math. Inf. Sci. (2013), in press.
[20] Inan, E. and Öztürk, M. A. *Fuzzy soft rings and fuzzy soft ideals*, Neural Comput. and Applic DOI 10.1007/s00521-011-0550-5.
[21] Jun, Y. B. and Park, C.H. *Applications of soft sets in ideal theory of BCK/BCI-algebras*, Inform. Sci. **178**, 2466-2475, 2008.
[22] Kong, Z., Gao, L.Q. and Wang, L.F. *Comment on a fuzzy soft set theoretic approach to decision making problems*, J. Comput. Appl. Math. **223**, 540-542, 2009.
[23] Maji, P. K., Roy, A. R. and Biswas, R. *An application of soft sets in desicion making problem*, Comput. Math. Appl. **44**, 1077-1083, 2002.
[24] Maji, P. K., Biswas, R. and Roy, A. R. *Fuzzy soft sets*, Journal of Fuzzy Mathematics, **203** (2), 589-602, 2001.
[25] Ming, P. B. and Ming, L. Y. *Fuzzy topology I.Neighbourhood structure of a fuzzy point and Moore-Smith convergence*, J. Math. Anal. Appl. **76,** 571-599, 1980.
[26] Molodtsov, D. *Soft set theory-First results*, Comput. Math. Appl. **37** (4/5), 19-31, 1999.
[27] Shabir, M. and Naz, M. *On soft topological spaces*, Comput. Math. Appl. **61**, 1786-1799, 2011.
[28] Roy, A.R. and Maji, P.K. *A fuzzy soft set theoretic approach to decision making problems*, J. Comput. Appl. Math. **203**, 412-418, 2007.
[29] Tanay, B. and Kandemir, M. B. *Topological structures of fuzzy soft sets*, Comput. Math. Appl. **61**, 412-418, 2011.
[30] Şimsekler, T. and Yüksel, S. *Fuzzy soft topological spaces*, Ann. Fuzzy Math. Inform. **5** (1), 87-96, 2013.
[31] Xiao, Z., Gong, K. and Zou, Y. *A combined forecasting approach based on fuzzy soft sets*, J. Comput. Appl. Math. **228**, 326-333, 2009.
[32] Zadeh, L. A. *Fuzzy sets*, Inform. and Control **8**, 338-353, 1965.
[33] Zorlutuna, İ., Akdağ, M., Min, W. K. and Atmaca, S. *Remarks on soft topological spaces*, Ann. Fuzzy Math. Inform. **3** (2), 171-185, 2011.

Cumhuriyet University, Department of Mathematics, 58140 Sivas / Turkey
*E-mail address*: seatmaca@cumhuriyet.edu.tr

Cumhuriyet University, Department of Mathematics, 58140 Sivas / Turkey
*E-mail address*: izorlu@cumhuriyet.edu.tr